

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/259268308>

# Long-term Outcome in Patients with Recurrent Malignant Glioma Treated with Perillyl Alcohol Inhalation

Article in *Anticancer research* · December 2013

Impact Factor: 1.83 · Source: PubMed

---

CITATIONS

9

---

READS

33

7 authors, including:



[Clovis Orlando da Fonseca](#)

Universidade Federal Fluminense

31 PUBLICATIONS 307 CITATIONS

SEE PROFILE



[Juliana de Saldanha da Gama Fischer](#)

Fundação Oswaldo Cruz

26 PUBLICATIONS 293 CITATIONS

SEE PROFILE



[José Alberto Landeiro](#)

Universidade Federal Fluminense

48 PUBLICATIONS 228 CITATIONS

SEE PROFILE



[Thereza Quirico-Santos](#)

Universidade Federal Fluminense

59 PUBLICATIONS 611 CITATIONS

SEE PROFILE

# ANTICANCER RESEARCH

International Journal of Cancer Research and Treatment

ISSN: 0250-7005 (print)  
ISSN: 1791-7530 (online)

*Editorial Office:* International Institute of Anticancer Research,  
DELINASIOS G.J. & CO G.P., Kapandriti, POB 22, Attiki 19014,  
Greece. Fax: 0030-22950-53389; Tel: 0030-22950-52945  
e-mail: journals@iiar-anticancer.org

Dear Sir/Madam:

Enclosed are the galley proofs of your article for ANTICANCER RESEARCH.

We would like to call your attention to the following:

1. Please read thoroughly, correct, and return the proofs to the Editorial Office **within 24 hours**.
2. Proofs should be returned preferably **by e-mail** or **fax**. Delays in the return of these proofs will necessitate the publication of your paper in a later issue of the journal.
3. Please read the entire manuscript carefully to verify that no changes in meaning have been introduced into the text through language improvements or editorial corrections.
4. Corrections should be limited to typographical errors.
5. Should you require reprints, PDF file, online open access, issues or special author rate subscriptions, please fill the attached reprint order form.
6. If you opt for online open access publication of your paper, it will instantly upon publication be available to read and reproduce free of charge.
7. Should you require information about your article (publication date, volume, page numbers, etc) please call: +30-22950-52945 or send an e-mail to journals@iiar-anticancer.org.
8. Please provide your complete address (not P.O.B.), telephone and fax numbers for the delivery of reprints and issues.
9. Please feel free to contact us with any queries that you may have (Tel./Fax: +30-22950-53389 or +30-22950-52945, e-mail: journals@iiar-anticancer.org).

Thank you for taking the time to study these guidelines.

I greatly appreciate your cooperation and your valuable contribution to this journal.

Yours sincerely,



*J.G. Delinasios*  
*Managing Editor*

Enclosures

## Long-term Outcome in Patients with Recurrent Malignant Glioma Treated with Perillyl Alcohol Inhalation

CLOVIS O DA FONSECA<sup>1</sup>, RAPHAEL M. TEIXEIRA<sup>2</sup>, JÚLIO CESAR T. SILVA<sup>3</sup>, JULIANA DA GAMA FISCHER<sup>4</sup>, OSÓRIO C. MEIRELLES<sup>5</sup>, JOSE ALBERTO LANDEIRO<sup>1</sup> and THEREZA QUIRICO-SANTOS<sup>2</sup>

<sup>1</sup>Department of General and Specialized Surgery, Antonio Pedro University Hospital, Fluminense Federal University, Rio de Janeiro, Brazil;

<sup>2</sup>Department of Cellular and Molecular Biology, Institute of Biology, Fluminense Federal University, Rio de Janeiro, Brazil;

<sup>3</sup>Department of Neurosurgery, Ipanema Federal Hospital, Rio de Janeiro, Brazil;

<sup>4</sup>Laboratory for Proteomics and Protein Engineering, Carlos Chagas Institute, Fiocruz/Paraná. Brazil;

<sup>5</sup>Rio de Janeiro Federal University, Rio de Janeiro, Brazil

**Abstract.** *Aim: This retrospective study aimed to evaluate long-term response and toxicity of recurrent malignant glioma patients to inhalation chemotherapy with perillyl alcohol (POH). Patients and Methods: The cohort included 117 men and 81 women with primary glioblastoma multiforme (GBM; n=154), grade III astrocytoma (AA; n=26) and anaplastic oligodendroglioma (AO; n=5). POH inhalation schedule 4-times daily started with 66.7 mg/dose; 266 mg/day and escalated up to 133.4 mg/dose; 533.6 mg/day. Clinical toxicity and overall survival following treatment were compared with tumor size; topography; extent of peritumoral edema and histological classification. Results: Adhesion to the protocol was high (>95%), POH (533.6 mg/daily) occasionally caused nose soreness but rarely nosebleed. Tumor size, peritumoral edema and oligodendroglial component influenced response to treatment. Conclusion: After 4 years under exclusive POH treatment, 19% patients still remain in clinical remission. Long-term POH inhalation chemotherapy is a safe and non-invasive strategy efficient for recurrent malignant glioma.*

Malignant gliomas are the most common primary intracranial tumors, and glioblastoma multiforme (GBM) accounts for 15% of all adult primary brain tumors (1). Despite aggressive multimodal treatment comprising of surgical resection, local radiotherapy (RT) and systemic chemotherapy the median

survival time after diagnosis is still in the range of 12 months, but population-based studies indicate an even shorter survival (2-3). Standard treatment includes maximal safe surgical resection, greater than 98% of the tumor volume to improve prognosis, but GBM tumors are poorly-delineated, with infiltrating tumor cells, intense angiogenesis and necrosis occupying more than 90% of the tumor mass. GBM recurrence occurs within 6.9 months after diagnosis, and all patients recur after initial therapy with new lesion appearing in the same tumor area (4), and very short (4 to 6 months) survival after first recurrence (5). For this reason, most GBM patients undergo standard-regimen combination of radiotherapy and adjuvant temozolomide (TMZ) chemotherapy (4).

TMZ is an orally-administered alkylating pro-drug capable to alkylate/methylate DNA by delivering a methyl group to purine DNA (O6-guanine; N7-guanine and N3-adenine) that damages DNA and triggers tumor cell death. DNA repair depends on the expression of O-6-methylguanine-DNA methyltransferase (MGMT) enzyme that diminishes the therapeutic efficacy of TMZ in tumors expressing this protein, or tolerated in mismatch repair-deficient (MMR-) tumors (4, 6). GBM initiation, recurrence and drug resistance is associated to modification of enzymes, non-coding RNAs and cell mutations in different cell types from the original glioma cell passing on mutations to oligodendrocyte or astrocyte precursor cells (4, 7). Besides, a relatively quiescent subset of endogenous glioma cells, albeit with stem cell properties, sustains long-term tumor growth through the production of highly proliferative cells (4, 8). Standard salvage therapies include various chemotherapeutic drugs, repeat resection, radiosurgery or interstitial implantation. Treatment decisions are complicated because additional RT may pose risk of cumulative toxicity, and options for chemotherapy may be limited by the development of resistance (5). To improve the clinical

*Correspondence to:* Thereza Quirico-Santos, Laboratory of Cellular Pathology, Institute of Biology, Fluminense Federal University, Niterói, RJ, 24020-141, Brazil. Tel: +55 2126292305, Fax: +55 2126292268, e-mail: tquirico@vm.uff.br

*Key Words:* Phase II trial, perillyl alcohol, intranasal administration, glioma, GBM, glioblastoma multiforme.

outcome in patients with malignant glioma, novel chemotherapeutic agents and effective new regimens are needed. Molecular alterations in neoplastic glioma often result from multiple mutations in proto-oncogenes and tumor suppressor genes (9) encoding for proteins that play important roles in signal transduction abnormalities of three major pathways: extracellular growth factors and their receptors EGF/EGFR, PDGF/PDGFR, and the signal transduction cascades Ras and AKT (10-11). Therapies targeting these specific molecules may result in killing tumor cells effectively while keeping normal cells intact. In this context, the search for new chemotherapeutic drugs has increased, especially for bioactive natural compounds (12).

Perillyl alcohol (POH) is a non-toxic naturally-occurring hydroxylated monoterpene with proved pre-clinical activity against various tumors in humans including breast, pancreas and lung carcinomas (13-15). As a cytotoxic agent, POH inhibits cell cycle, up-regulates the pro-apoptotic protein Bax (15-17) and also TMZ-resistant and TMZ-sensitive glioma cells, independently of O6-methylguanine-DNA methyltransferase (MGMT) expression (18). POH cytotoxicity upon glioma cell lines resulted from the effect on endoplasmic reticulum (ER) stress pathway, as shown by increased expression of glucose-regulated protein-78 (GRP78), transcription factor-3 (ATF3) activation, and C/EBP-homologous protein (CHOP), and also by arresting survival pathways, such as mTOR and Ras (18-19).

The brain tissue is one of the least accessible organs for delivery of active pharmacological compounds (20). Our group has pioneered on the use of intranasal POH for the treatment of patients with recurrent malignant glioma (Phase I/II) (21-22). Intranasal administration allows POH to cross the blood-brain barrier (BBB) and reaches the central nervous system (CNS) thus eliminating the need for systemic delivery while reducing side-effects. We present data (Phase II study) showing that long-term administration of POH as a single agent by inhalation was an effective therapeutic strategy capable to sustain long-term regression (>4 years survival) of recurrent glioma without clinical systemic side-effects.

## Patients and Methods

This clinical trial was approved by the Hospital Medical Research Ethics Committee and the Brazilian Ministry of Health (CONEP 9681 no. 25000.009267/2004-25, July 12, 2004) complies with the principles laid down in the Declaration of Helsinki. Informed consent was obtained from each patient and the next-of-kin before beginning the study.

*Patient selection.* This prospective study was carried-out from July 2006 to July 2012 with patients attending the out-patient Neurosurgical Unit in the Antonio Pedro University Hospital and included in the Phase I/II clinical trial to assess the efficacy of intranasal administration of the monoterpene POH. The cohort included after appropriate informed consent of 198 adult patients

(random selection) with recurrent malignant glioma and under symptomatic treatment after failing response to current standard treatment that included surgery, and/or radiation, and multimodal chemotherapy specific for glioblastoma. Diagnosis and histological classification of malignant glioma was based upon the WHO criteria (23). Eligibility criteria included patients older than 18 years with recurrent glioblastoma with at least two relapses, measurable contrast-enhancing tumor image on magnetic resonance, Karnofsky index  $\geq 70\%$ , adequate hematological clinic laboratory-based measures, stable heart rhythm and no clinical evidence of congestive heart failure or unstable angina. Exclusion criteria included pregnancy, hematological malignancy, occurrence of seizures, recurrence within the last 96 h before inclusion; concomitant infectious or inflammatory processes; acute cerebrovascular or hemorrhagic events. Classification according to WHO was: 155 (78%) glioblastoma IV (GBM); 27 (14%) anaplastic astrocytoma (AA) and 16 (8%) anaplastic oligodendroglioma (AO) patients, who were followed by MRI and clinical evaluation every 3 months. Adverse events were graded according to Common Terminology Criteria for Adverse Events–Version 4.0 (CTCAE). At the moment of the study, none were under radiation therapy or chemotherapy for more than 4 weeks. All patients were under palliative symptomatic treatment because they had failed current standard treatment for glioma recurrence.

A correlation of initial symptoms and clinical presentation with topographical localization (lobar or deep) and size of the tumor in the brain tissue was done based on brain MRI image. Lobar tumors were defined as any tumoral lesion away from the basal ganglia. All tumors limited to or involving the basal ganglia were classified as deep. Tumor size was measured on axial contrast-enhanced scans using the scale of the largest perpendicular diameters of the enhancing lesion. The extent of peritumoral edema (PTBE), mass effect and midline shift (cm) were evaluated at initial diagnosis and during the course of treatment with monoterpene POH. The area occupied by peritumoral edema was determined on the same axial slices used for tumor size measurements by subtracting tumor from edema diameters. The comparison was kept constant with same type of axial imaging (CT or MRI).

*Drug administration and dose escalation.* Perillyl alcohol was formulated for intranasal delivery and the preparation supplied by the University Pharmacy was according to the following patents: US Patent Application 20040087651 May 6, 2004, and BR Application Number 0107262-5 December 17, 2001. Perillyl alcohol (Sigma Chem. Co., St Louis, MI, USA) 0.3% v/v POH (55 mg) was administered by inhalation 4 times daily. All patients received POH 4 times daily by intranasal (inhalation) delivery from initial dose (66.7 /dose; totaling 266.8 mg/daily), and escalation 133.4 mg/dose, totaling 533.6 mg/daily.

*Statistical analysis.* Statistical analysis was carried out using Kaplan–Meier curves, log rank tests and univariate and multivariate Cox regression models. Association of variables with survival was assessed with Kaplan–Meier curves and log rank tests, and significance of continuous variables with univariate Cox regression models.

## Results

Demographic characteristic of patients are specified in Table I. Patients were stratified in two groups: recurrent primary GBM and recurrent secondary GBM derived from low-grade

Table I. Patient characteristics (n=198).

	GBM	AA	AO
Histology	155 (78%)	27 (27%)	16 (8%)
Tumor type			
recurrent primary GBM	144 (92%)		
recurrent secondary GBM	11 (8%)		
Age, years; mean (range)	51 (38-64)	52 (41-64)	49 (28-69)
Gender			
male	92 (59%)	16 (59%)	9 (56%)
female	63 (41%)	11 (41%)	7 (44%)
Initial predominant symptom (GBM, AA, AO)			
headache		53%	
seizures		25%	
focal neurological deficit		52%	
short-term memory loss		11%	
temporary confusion		14%	
behavioral changes		7%	
Site of tumor			
lobar	122 (78%)	22 (80%)	16 (100%)
deep	33 (21%)	5 (20%)	none
Treatment of the initial tumor			
gross total resection	78%	77%	100%
subtotal resection or biopsy	22%	23%	0%
radiotherapy	100%	100%	100%
chemotherapy	100%	100%	100%
PFS with POH intranasally (month 6th)	48%	60%	66%
Edema on initial tumor (MRI)			
none	5 (9.6%)	1 (10%)	3 (60%)
<5 cm	24 (46.2%)	7 (70%)	2 (40%)
>5 cm	28 (53.8%)	2 (20%)	none
Edema on recurrent tumor (MRI)			
<5 cm	19 (36.5%)	6 (60%)	5(100%)
>5 cm	33 (63.4%)	4 (40%)	1
Midline shift on initial tumor (MRI)			
none	8 (15.3%)	2 (20%)	3 (60%)
<1 cm	19 (36.5%)	6 (60%)	2 (40%)
>1 cm	25 (48.7%)	2 (20%)	none
Midline shift on recurrent tumor (MRI)			
<1 cm	18 (35.5%)	4 (40%)	3 (60%)
>1 cm	34 (64.5%)	6 (60%)	2 (40%)

glioma. Gross total resection was performed in 103 patients, whereas 95 patients underwent subtotal resection or stereotactic biopsy. According to CTCAE, 5 patients had nasal aching, and 2 patients had epistaxis after prolonged use with 133.4 mg/dose: 533.6 mg/daily POH. After topic treatment, POH dose was reduced (266.8 mg/daily) and patients improved clinical condition without any further complaint. There was no evidence of local or systemic infection, gastrointestinal toxicity and hematological side-effects, even in patients with prolonged treatment with intranasal POH for more than 4 years.

The most frequent complaint at onset was intense headache (53%), but patients also reported focal neurological signal - hemiparesis (29%); convulsive seizures (25%);

nausea and mental confusion (14%); visual dizziness and lack of memory (11%); behavior alteration, deficit of language and cognition (7%), sleepiness (4%) before diagnostic confirmation of malignant brain tumor. Prevalent complaints at tumor recurrence were neurological deficits (51%); intense headache (43%) and seizures in 24% (Table I).

Long-term treatment (up to 4 years) with daily intranasal POH administration as single-chemotherapy drug, stabilized (n=3) and improved clinical condition (n=2) of patients with grade II astrocytoma (Figure 1). However due to small number of AA (n=27) patients included, further studies are required to determine the effect of treatment on the natural history of the disease. AA patients consistently showed




Figure 1. Effect of intranasal POH treatment in patients with astrocytoma grade II. Representative MRIs of AA-II patient (A) show marked reduction of tumor size after 2 years and 3 years of POH treatment in comparison with first image obtained before treatment. (B) Mild reduction of tumor size after 2 and 3 years of POH treatment in comparison with first image obtained before treatment.




Figure 2. Overall survival of glioma patients included in Phase I/III clinical trial. Kaplan-Meier curves show the influence of long-term treatment without discontinuation of intranasal POH administration. Comparison of primary GBM (A) and secondary GBM ( $p < 0.0001$ ); (B) anaplastic astrocytoma (AA) and anaplastic oligodendroglioma ( $p = 0.0199$ ). Line of tendency indicates survival rate of patients.

evident reduction of tumor size in comparison with MRI image taken before patient inclusion in POH treatment. However, it is important to establish a relation between the effectiveness of treatment with topography and histo-




Figure 3. Effect of intranasal POH treatment in recurrent oligoastrocytoma. A patient with recurrent oligoastrocytoma and extensive peritumoral edema failed to respond to POH treatment. MRI image obtained before (A) and 4 months after POH treatment (B). Good responder patient showed in comparison with initial MRI (C) a marked reduction (D) in the tumor size 4 years after POH treatment. HE staining (E) show increased glomeruloid microvascular and glial cell proliferation with atipia, hyperchromatism and nuclear pleomorphism. Brain biopsy fragment (F) obtained 4 years after intranasal POH treatment during intracranial drainage of hematoma show apparently normal characteristics of glial and neuronal cellularity.

molecular characteristics of the tumor, because even patients with dull response to POH treatment showed albeit reduction of tumor size after intranasal POH treatment.

A 6-month progression-free survival rate including partial responses and stable disease of glioma patients according to the histological diagnosis was GBM: 48%; AA: 60% and AO: 67%. Survival rate after 24 months of POH treatment was 6.2 % for primary GBM, 63% for secondary GBM comprising 15% for AA and 56% for AO. Kaplan-Meier




Figure 4. Effect of intranasal POH treatment in recurrent GBM. Representative brain image (MRI) of a patient with recurrent primary GBM. Decrease in the tumor size between the initial MRI (A, B) and 6 months after treatment (C, D).

curves (Figure 2) showed that patients with secondary GBM had high survival rates in comparison with primary GBM ( $p < 0.0001$ ) (Figure 2A) evidenced by a sharp decline of tendency line. Patients with anaplastic astrocytoma (AA) showed a significant ( $p = 0.0199$ ) decline of tendency line compared with anaplastic oligodendroglioma (Figure 2B), that conversely had prolonged survival rate further evidenced by a mild and steady decline of tendency line.

Lobar location was present in 80.8% of glioma patients (Table I), and as previously reported (24) it was also observed a correlation of tumor topography with therapeutic response to intranasal administration of POH. Patients with tumoral lesion in the basal ganglia survived longer than those with tumors at a lobar localization. Likewise, presence of peritumoral edema, midline shift and requirement for continued use of steroids also influenced morbidity and tumor invasive recurrence. Furthermore, maintenance of extensive brain edema ( $> 5$  cm) and midline shift ( $> 1$  cm) since the initial (onset) MRI scan despite regression of tumor size, determined a poor response to POH treatment and shorter survival than patients with moderate edema (Figure 3). Nonetheless, patients either with oligodendroglioma or anaplastic oligodendroglioma (AO) showed good response to intranasal POH administration (Figure 2). Patients with AO or oligodendroglioma but with peritumoral edema greater




Figure 5. Tumor recurrence after treatment discontinuation. Decrease in the tumor image size of a patient with secondary GBM at the time of inclusion in POH trial (A); image obtained 3 years after continuous treatment with intranasal POH (B); and image (C) obtained three months after patient have quit treatment.

than 5 cm also had poor response to POH (Figure 3). Good responders to POH treatment included patients with prolonged survival that were not using steroids for a long period of time. Conversely, maintenance of high steroid dosage besides persistence of large peritumoral edema was often associated with negative response to POH treatment, and morbidity with recurrent infections, and deep venous and carotid artery thrombosis.

Fully-adhesion to POH treatment was of utmost importance because discontinuation (Figure 5) caused tumor recurrence even after early MRI image showing marked reduction of the tumoral lesion. Prolonged POH inhalation chemotherapy did not cause cumulative toxicity, and neither altered clinical chemistry (hepatic, renal, lung) and hematological parameters. Glioma patients (19%) with good response to POH treatment and improvement of clinico-neurological status without evidence of MRI tumor recurrence had their POH dosage reduced to the lower concentration.

## Discussion

Standard chemotherapeutic regimens for treatment of brain tumors are frequently inefficient due to the inability of drugs to reach and maintain effective concentrations within the brain tissue for an adequate length of time (20). The role of olfactory epithelium as a gateway for substances entering the CNS have lead us to develop a novel and ground-breaking methodology for POH delivery to treat patients with recurrent malignant brain tumors (21-22). Intranasal delivery is a non-invasive method that allows rapid access of drugs and hormones to the CNS, bypassing the BBB, minimizing systemic exposure and dramatically reducing adverse side-effects (25-26). The present study was conducted to assess the comparative overall survival, progression-free survival, response rate, and safety of long-term POH inhalation chemotherapy among patients with recurrent malignant glioma.

A factor that greatly influenced the overall survival and response of glioma patients to long-term intranasal POH treatment was the presence of peritumoral edema that contributed to clinical symptoms (intense headache, dizziness, focal neurological deficit and seizures) and morbidity, and eventually favored glioma cell invasion to other brain structures. Persistence of peritumoral edema associated with midline shift and the need for continued use of steroids was also an indication of unfavorable prognosis. Even patients with oligodendroglioma or anaplastic oligodendroglioma that generally were good responders to intranasal POH administration had poor prognosis and dull response to treatment whenever peritumoral edema was greater than 5 cm.

Although malignant gliomas do not produce metastases, by the time of diagnosis and treatment, glioma cells have already actively-invaded the surrounding brain parenchyma producing undetectable tumor niches which unable effective surgical resection (27-28). In fact, studies using spectroscopy imaging showed the presence of neoplastic cell activity in areas apparently healthy without any macroscopic evidence of tumoral activity (29-30). Patients with worse prognosis and poor response to intranasal POH treatment had tumoral lesion in the supratentorial region which presents a dense microvascular network towards the brainstem. The tumoral microenvironment is rich in growth factors and tissue proteases that diffuse through the peritumoral stroma and activate intracellular pathways, thereby promoting extensive neovascularization and generation of epigenetic events that promotes new oncogenic mutations, and lost-in-function of different tumor suppressor genes (27, 31). Indeed, abnormal methylation of CpG promoter regions of several genes has been recognized as an important mechanism of epigenetic gene silencing, contributing to changes in gene expression and resistance to alkylating drugs (32). For development of new therapeutic approaches, it is important to determine major molecular pathways responsible for proliferation, invasion, angiogenesis, and anaplastic transformation of glioma cells. We demonstrated that intranasal administration of a p21-Ras lipophilic inhibitor that easily crosses the blood-brain barrier is a safe and noninvasive strategy capable to prolong overall survival of patients with recurrent malignant glioma considered at terminal stage. In conclusion, (1) long-term POH inhalation was a safe and noninvasive therapeutic strategy for malignant gliomas; (2) long-term POH inhalation consistently improved survival of patients with grade III and grade IV anaplastic astrocytoma with oligodendroglioma component; (3) patients with secondary GBM had a better response to POH inhalation than patients with primary GBM; (4) 19% recurrent malignant glioma patients still remain in clinical remission after 4 years under exclusive POH inhalation treatment; (5) long-term POH inhalation did not cause any evident clinical and /or

laboratory-based toxicity. We can envisage in a near future the synthesis of biological active hybrid molecules containing POH as a carrier conjugated to drugs specifically targeting critical regulators of cell proliferation, as a promising antitumor therapeutic strategy successfully employed to treat brain tumors.

### Acknowledgments

This study was supported in part by grants from National Council Research (MCT/CNPq/CT-Saude 401943/2010-0; CNPq/Universal 481059/2011-3), Rio de Janeiro Research Foundation (FAPERJ: E-26/ 110.329/2011; E-26/110.948/2013), University Research Funding (FOPESQ 2013), and Euclides da Cunha Foundation/FEC-UFF.

### References

- 1 Adamson DC, Rasheed BA, McLendon RE and Bigner DD: Central nervous system. *Cancer Biomark* 9: 193-210, 2010.
- 2 Ohgaki H, Dessen P, Jourde B, Horstmann S, Nishikawa T, DiPatre P, Burkhard C, Schuler D, Probst-Hensch N, Maiorka P, Baeza N, Pisani P, Yonekawa Y, Yasargil M, Lutolf U and Kleihues P: Genetic pathways to glioblastoma: a population-based study. *Cancer Res* 64: 6892-6899, 2004.
- 3 Hoelzinger DB, Demuth T and Berens ME: Autocrine Factors That Sustain Glioma Invasion and Paracrine Biology in the Brain Microenvironment. *Journal of the National Cancer Institute* 99: 1583-1593, 2007.
- 4 Clarke J, Penas C, Pastori C, Komotar RJ, Bregy A, Shah AH, Wahlestedt C and Ayad NG: Epigenetic pathways and glioblastoma treatment. *Epigenetics* 8: 785-795, 2013.
- 5 Minniti G, Scaringi C, De Sanctis V, Lanzetta G, Falco T, Di Stefano D, Esposito V and Enrici RM: Hypofractionated stereotactic radiotherapy and continuous low-dose temozolomide in patients with recurrent or progressive malignant gliomas. *J Neurooncol* 111: 187-194, 2012.
- 6 Zhang J, Stevens MF and Bradshaw TD: Temozolomide: mechanisms of action, repair and resistance. *Curr Mol Pharmacol* 5: 102-114, 2012.
- 7 Verhaak RGW, Hoadley KA, Purdom E, Wang V, Qi Y, Wilkerson MD, Miller CR, Ding L, Golub T, Mesirov JP, Alexe G, Lawrence M, O'Kelly M, Tamayo P, Weir BA, Gabriel S, Winckler W, Gupta S, Jakkula L, Feiler HS, Hodgson JG, James CD, Sarkaria JN, Brennan C, Kahn A, Spellman PT, Wilson RK, Speed TP, Gray JW, Meyerson M, Getz G, Perou CM and Hayes DN: Integrated Genomic Analysis Identifies Clinically Relevant Subtypes of Glioblastoma Characterized by Abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell* 17: 98-110, 2010.
- 8 Chen J, Li Y, Yu TS, McKay RM, Burns DK, Kernie SG and Parada LF: A restricted cell population propagates glioblastoma growth after chemotherapy. *Nature* 488: 522-526, 2012.
- 9 Nigro JM, Misra A, Zhang L, Smirnov I, Colman H, Griffin C, Ozburn N, Chen M, Pan E, Koul D, Yung WK, Feuerstein BG and Aldape KD: Integrated array-comparative genomic hybridization and expression array profiles identify clinically relevant molecular subtypes of glioblastoma. *Cancer Res* 65: 1678-1686, 2005.

- 10 Lassman AB and Holland EC: Incorporating molecular tools into clinical trials and treatment for gliomas? *Curr Op Neurol* 20: 708-711, 2007.
- 11 Karp JE and Lancet JE: Development of farnesyltransferase inhibitors for clinical cancer therapy: focus on hematologic malignancies. *Cancer Invest* 25: 484-494, 2007.
- 12 Duelund L, Amiot A, Fillon A and Mouritsen OG: Influence of the Active Compounds of *Perilla frutescens* Leaves on Lipid Membranes. *Journal of Natural Products* 75: 160-166, 2012.
- 13 Wiseman DA, Werner SR and Crowell PL: Cell Cycle Arrest by the Isoprenoids Perillyl Alcohol, Geraniol, and Farnesol Is Mediated by p21<sup>Cip1</sup> and p27<sup>Kip1</sup> in Human Pancreatic Adenocarcinoma Cells. *Journal of Pharmacology and Experimental Therapeutics* 320: 1163-1170, 2007.
- 14 Bailey HH, Attia S, Love RR, Fass T, Chappell R, Tutsch K, Harris L, Jumonville A, Hansen R, Shapiro GR and Stewart JA: Phase II trial of daily oral perillyl alcohol (NSC 641066) in treatment-refractory metastatic breast cancer. *Cancer Chemother Pharmacol* 62: 149-157, 2008.
- 15 Yeruva L, Pierre KJ, Elegbede A, Wang RC and Carper SW: Perillyl alcohol and perillic acid induced cell cycle arrest and apoptosis in non small cell lung cancer cells. *Cancer Letters* 257: 216-226, 2007.
- 16 Holstein SA and Hohl RJ: Monoterpene regulation of Ras and Ras-related protein expression. *J Lipid Res* 44: 1209-1215, 2003.
- 17 Mo H and Elson CE: Studies of the isoprenoid-mediated inhibition of mevalonate synthesis applied to cancer chemotherapy and chemoprevention. *Exp Biol Med (Maywood)* 229: 567-585, 2004.
- 18 Cho HY, Wang W, Jhaveri N, Torres S, Tseng J, Leong MN, Lee DJ, Goldkorn A, Xu T, Petasis NA, Louie SG, Schonthal AH, Hofman FM and Chen TC: Perillyl alcohol for the treatment of temozolomide-resistant gliomas. *Mol Cancer Ther* 11: 2462-2472, 2012.
- 19 Sundin T, Peffley DM, Gauthier D and Hentosh P: The isoprenoid perillyl alcohol inhibits telomerase activity in prostate cancer cells. *Biochimie* 94: 2639-2648, 2012.
- 20 Caruso G, Caffo M, Alafaci C, Raudino G, Cafarella D, Lucerna S, Salpietro FM and Tomasello F: Could nanoparticle systems have a role in the treatment of cerebral gliomas? *Nanomedicine* 7: 744-752, 2011.
- 21 Da Fonseca CO, Schwartsmann G, Fischer J, Nagel J, Futuro D, Quirico-Santos T and Gattass CR: Preliminary results from a phase I/II study of perillyl alcohol intranasal administration in adults with recurrent malignant gliomas. *Surg Neurol* 70: 259-266, 2008.
- 22 Da Fonseca CO, Simao M, Lins IR, Caetano RO, Futuro D and Quirico-Santos T: Efficacy of monoterpene perillyl alcohol upon survival rate of patients with recurrent glioblastoma. *J Cancer Res Clin Oncol* 137: 287-293, 2011.
- 23 Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, Scheithauer BW and Kleihues P: The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 114: 97-109, 2007.
- 24 Da Fonseca CO, Silva JT, Lins IR, Simao M, Arnobio A, Futuro D and Quirico-Santos T: Correlation of tumor topography and peritumoral edema of recurrent malignant gliomas with therapeutic response to intranasal administration of perillyl alcohol. *Invest New Drugs* 27: 557-564, 2009.
- 25 Dhuria SV, Hanson LR and Frey WH: Intranasal delivery to the central nervous system: Mechanisms and experimental considerations. *Journal of Pharmaceutical Sciences* 99: 1654-1673, 2010.
- 26 Pardeshi CV and Belgamwar VS: Direct nose to brain drug delivery via integrated nerve pathways bypassing the blood-brain barrier: an excellent platform for brain targeting. *Expert Opinion on Drug Delivery* 10: 957-972, 2013.
- 27 De Palma M and Hanahan D: The biology of personalized cancer medicine: Facing individual complexities underlying hallmark capabilities. *Molecular Oncology* 6: 111-127, 2012.
- 28 Bellail AC, Hunter SB, Brat DJ, Tan C and Van Meir EG: Microregional extracellular matrix heterogeneity in brain modulates glioma cell invasion. *The International Journal of Biochemistry & Cell Biology* 36: 1046-1069, 2004.
- 29 Moller-Hartmann W, Herminghaus S, Krings T, Marquardt G, Lanfermann H, Pilatus U and Zanella F: Clinical application of proton magnetic resonance spectroscopy in the diagnosis of intracranial mass lesions. *Neuroradiology* 44: 371-381, 2002.
- 30 Server A, Josefsen R, Kulle B, Maehlen J, Schellhorn T, Gadmar O, Kumar T, Haakonsen M, Langberg CW and Nakstad PH: Proton magnetic resonance spectroscopy in the distinction of high-grade cerebral gliomas from single metastatic brain tumors. *Acta Radiol* 51: 316-325, 2010.
- 31 Esteller M: Epigenetics provides a new generation of oncogenes and tumour-suppressor genes. *Br J Cancer* 94: 179-183, 2006.
- 32 Di Vinci A, Casciano I, Marasco E, Banelli B, Ravetti GL, Borzi L, Brigati C, Forlani A, Dorcaratto A, Allemanni G, Zona G, Spaziante R, Gohlke H, Gardin G, Merlo DF, Mantovani V and Romani M: Quantitative methylation analysis of HOXA3, 7, 9, and 10 genes in glioma: association with tumor WHO grade and clinical outcome. *J Cancer Res Clin Oncol* 138: 35-47, 2012.

*Received October 28, 2013*

*Revised November 6, 2013*

*Accepted November 8, 2013*

# ANTICANCER RESEARCH

International Journal of Cancer Research and Treatment

ISSN: 0250-7005 (print) ISSN: 1791-7530 (online)

November 8, 2013

Dr. Thereza Quirico-Santos

Re: Your manuscript No. **16784-D** entitled «**Long-term Outcome in Patients...**»

Dear Dr

Referring to your above manuscript for publication in AR, please allow us to use this form letter in reply:

1. *Referee's recommendations:*

- Urgent to be published immediately.
- Accepted in the presented form.
- Accepted with minor changes.
- Accepted with grammatical or language corrections.
- Remarks:

2. *Excess page charges.*

- Your article has approx. **7** printed pages and is in excess of the allotted number by approx. **3** printed pages. The charges are EURO € **180** per excess page, totalling EURO € **540**  
We ask you to confirm acceptance of these charges.
- Your article includes **1** pages with color figures. The charges are EURO € **650** per color page, totalling EURO € **650**  
**TOTAL charges are EURO € 892.50 (-25% Discount)**
- Our invoice will be sent by e-mail to the corresponding author.

3.  Your article will appear in Volume **33**, Issue No. **12**, **2013**

4.  Please order your reprints, pdf or online open access, now. This will facilitate our prompt planning of future issues and rapid publication of your article. Reprints will be delivered by rapid one-day delivery within one month from publication.

We would appreciate your prompt reply.

With many thanks,

Yours sincerely,



J.G. Delinasios

Managing Editor

EDITORIAL OFFICE: INTERNATIONAL INSTITUTE OF ANTICANCER RESEARCH  
DELINASIOS G.J. & CO G.P., Kapandriti, P.O.B. 22, Attiki 19014, Greece. Tel.: 0030-22950-52945;  
Tel & Fax:0030-22950-53389; e-mail: journals@iiar-anticancer.org

# ANTICANCER RESEARCH

## International Journal of Cancer Research and Treatment

*Editorial Office:* International Institute of Anticancer Research,  
DELINASIOS G.J. & CO G.P., Kapandriti, P.O.B. 22, Attiki 19014, Greece  
Fax: +30-22950-53389; Tel: +30-22950-52945; e-mail: journals@iiar-anticancer.org

ISSN: 0250-7005 (print)  
ISSN: 1791-7530 (online)

Please type or print the requested information on the reprint order form and return it to the Editorial Office by fax or e-mail.

Fees for reprints, PDF file, online open access and subscriptions must be paid for in advance.

If your paper is subject to charges for excess pages or color plates, please add these charges to the payment for reprints. The reprints are not to be sold.

### PRICE LIST FOR REPRINTS WITHOUT COVER

Page length	Online		Number of copies requested (prices are in Euro)									
	Open Access Fee (Euro)	PDF File Fee (Euro)	50	100	200	300	400	500	1000	1500	2000	3000
1-4pp	400	175	140	285	337	388	453	504	851	1135	1470	2038
5-8	600	225	150	388	453	530	595	672	1083	1445	1832	2554
9-12	700	277	160	504	569	659	737	827	1341	1780	2219	3096
13-16	800	354	180	659	737	840	943	1046	1625	2141	2657	3676
17-20	900	419	200	788	879	982	1098	1227	1883	2451	3044	4244

*For reprints with cover: Please add EURO 120.00 per 100 copies.*

*Postage: Please add 5% on the reprint prices.*

## Reprint Order Form

Of my paper No. **16784-D** comprising **7** printed pages, entitled «**Long-term Outcome in Patients...**» accepted for publication in ANTICANCER RESEARCH Vol. **33** No. **12**

- I require a total of \_\_\_\_\_ copies at EURO:
- I do not require reprints.
- Please send me a PDF file of the article at EURO:
- Please provide Online Open Access of the article at the Stanford University Highwire Press website immediately upon publication at EURO:
- Please send me a copy of the issue containing my paper at EURO 45.00.
- Please enter my personal subscription to ANTICANCER RESEARCH at the special Author's price of EURO 390.00 ( print;  online) ( Year: 2013).
- A check for the above amounts payable to DELINASIOS G.J. & CO G.P., is enclosed.
- Please send an invoice to (Billing Name and Address):  
VAT number (For EC countries):  
Name:  
Address: \_\_\_\_\_ Signature:  
Country: \_\_\_\_\_ City:  
Postal code: \_\_\_\_\_ e-mail:  
Tel: \_\_\_\_\_ Fax:
- Please send reprints to (Complete Address and Tel. no.):